PNAS Online



HOME HELP FIEDBACK SUBSCRIPTIONS ARCHIVE SEARCH

Institution: PC43PT0608040 Sign In as Member / Individual

Supporting Information for Szyperski et al. (2002) Proc. Natl. Acad. Sci. USA 99 (12), 8009–8014. (10.1073/pnas.122224599).

▶ Full Text of this Article

Supporting Figure 13

Fig. 13. Sequential resonance assignment based on 3D $\underline{H}^{a/b}\underline{C}^{a/b}(CO)NHN / 3D \underline{H}^{a/b}\underline{C}^{a/b}COHA$ combined with 3D HNN< $\underline{CA,CO}$ > (see Fig. 2). The amino acid residue on which the NMR signal was detected is indicated at the bottom of the strips. (A) Matching of $w_1(^{13}C^{a/b}/^{1}H^{a/b})$ peak patterns in 3D $\underline{H}^{a/b}\underline{C}^{a/b}COHA$ (strips labeled with "a") and 3D $\underline{H}^{a/b}\underline{C}^{a/b}(CO)NHN$ ("b") yields putative intraresidue $^1H^{a/b}/^{13}C^{a/b}$ COHA the $^{13}C=O$ correlations: on the strips taken from 3D $\underline{H}^{a/b}\underline{C}^{a/b}COHA$ the $^{13}C=O$ chemical shift is indicated. (B) 3D HNN< $\underline{CA,CO}$ > yields the sequential $^{13}C=O^{-13}C^a$ correlations: the carbonyl chemical shifts have to match those shown in A, and are indicated on the left of the figure. Proton and carbon chemical shifts are given in ppm and relative to 2,2-dimethyl-2-silapentane-5-sulfonate (DSS).

Full Text of this Article

HOME HELP PEIDBACK SUBSCRIPTIONS ARCHIVE SEARCH

Copyright © by the National Academy of Sciences